

Nucleon Form Factors of the Isovector Axial-Vector Current

Situation of Experiments and Theory

M. R. Schindler and S. Scherer

Institut für Kernphysik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

Received: date / Revised version: date

Abstract. The theoretical and experimental status of the isovector axial-vector current form factors $G_A(q^2)$ and $G_P(q^2)$ of the nucleon is reviewed. We also describe a new calculation of these form factors in manifestly Lorentz-invariant chiral perturbation theory (ChPT) with the inclusion of axial-vector mesons as explicit degrees of freedom.

PACS. 11.40.-q Currents and their properties – 12.39.Fe Chiral Lagrangians

1 Introduction

The structure of the nucleon is encoded in several form factors. For example, the electromagnetic Dirac and Pauli form factors $F_1(q^2)$ and $F_2(q^2)$, or equivalently the electric and magnetic Sachs form factors $G_E(q^2)$ and $G_M(q^2)$, parameterize the matrix elements of the electromagnetic current operator and are well-known over a wide region of momentum transfer squared q^2 . For recent reviews on the experimental status of the electromagnetic form factors see, e. g., [1, 2, 3]. In contrast to the electromagnetic case, the three form factors of the isovector axial-vector current, $G_A(q^2)$, $G_P(q^2)$, and $G_T(q^2)$, are not as well-known. They encode the structure of the matrix elements of the isovector axial-vector current $A^{\mu,a}(x)$ which, in the SU(2) case, is given by

$$A^{\mu,a}(x) \equiv \bar{q}(x) \gamma^\mu \gamma_5 \frac{\tau^a}{2} q(x), \quad q = \begin{pmatrix} u \\ d \end{pmatrix}, \quad a = 1, 2, 3. \quad (1)$$

$A^{\mu,a}(x)$ is a Hermitian operator that transforms as an isovector under isospin transformations and as an axial vector under Lorentz transformations. The corresponding matrix element between initial and final nucleon states is parameterized as

$$\begin{aligned} \langle N(p') | A^{\mu,a}(0) | N(p) \rangle \\ = \bar{u}(p') \left[\gamma^\mu \gamma_5 G_A(q^2) + \frac{q^\mu}{2m_N} \gamma_5 G_P(q^2) \right. \\ \left. + i \frac{\sigma^{\mu\nu} q_\nu}{2m_N} \gamma_5 G_T(q^2) \right] \frac{\tau^a}{2} u(p), \end{aligned} \quad (2)$$

where $q = p' - p$ and m_N denotes the nucleon mass. $G_A(t)$ is called the axial form factor, $G_P(t)$ is the induced pseudoscalar form factor, and $G_T(t)$ denotes the induced pseudotensorial form factor. For the isospin-symmetric case of

equal up and down quark masses, $m_u = m_d$, the strong interactions are invariant under $\mathcal{G}$ conjugation, which is a combination of charge conjugation $\mathcal{C}$ and a rotation by π about the 2 axis in isospin space (charge symmetry operation),

$$\mathcal{G} = \mathcal{C} \exp(i\pi I_2). \quad (3)$$

As a consequence the pseudotensorial form factor $G_T(q^2)$ vanishes in the limit of perfect isospin symmetry. So far only upper limits have been established on the size of G_T [4]. We will neglect G_T in the following discussion. From the Hermiticity of the current operator one infers that G_A and G_P are real functions for space-like momentum transfer squared ($q^2 < 0$).

In the Breit system, where $\mathbf{p} = -\frac{1}{2}\mathbf{q} = -\mathbf{p}'$ and $q_0 = 0$, the zeroth component of the matrix element of Eq. (2) vanishes,

$$\langle N(\frac{\mathbf{q}}{2}) | A^0(0) | N(-\frac{\mathbf{q}}{2}) \rangle = 0, \quad (4)$$

while the spatial components can be written as¹

$$\begin{aligned} \langle N(\frac{\mathbf{q}}{2}) | \mathbf{A}(0) | N(-\frac{\mathbf{q}}{2}) \rangle = \\ [2E \boldsymbol{\sigma}_\perp G_A(-\mathbf{q}^2) + 2m_N \boldsymbol{\sigma}_\parallel D(-\mathbf{q}^2)] \frac{\boldsymbol{\tau}^a}{2}, \end{aligned}$$

where $E = \sqrt{m_N^2 + \mathbf{q}^2/4}$, $\boldsymbol{\sigma}_\parallel = \boldsymbol{\sigma} \cdot \hat{\mathbf{q}}$, $\boldsymbol{\sigma}_\perp = \boldsymbol{\sigma} - \boldsymbol{\sigma}_\parallel$, and

$$D(q^2) = G_A(q^2) + \frac{q^2}{4m_N^2} G_P(q^2).$$

We have thus separated the spatial components of the matrix element into its transverse and longitudinal parts. As we will see below, the longitudinal part vanishes in the chiral limit.

For recent reviews on the form factors of the isovector axial-vector current see [5, 6].

¹ We adopt the normalization $\bar{u}u = 2m_N$.

2 Axial form factor $G_A(q^2)$

$G_A(q^2)$ is the best known of the isovector axial-vector current form factors. Its value at zero momentum transfer squared is defined as the axial-vector coupling constant g_A ,

$$G_A(0) = g_A, \quad (5)$$

which has been determined from neutron beta decay to be [7]

$$G_A(q^2 \approx 0) = g_A = 1.2695 \pm 0.0029. \quad (6)$$

There are two distinct experimental ways of determining the q^2 dependence of G_A . The first is (quasi)elastic (anti)neutrino scattering (referred to as simply neutrino scattering in the following). For the analysis of experimental data, $G_A(q^2)$ is conventionally parameterized using a dipole form as

$$G_A(q^2) = \frac{g_A}{(1 - \frac{q^2}{M_A^2})^2}, \quad (7)$$

where M_A is the so-called axial mass. The global average for the axial mass extracted from neutrino scattering experiments given in [5] is

$$M_A = (1.026 \pm 0.021) \text{ GeV}, \quad (8)$$

whereas a recent analysis [8] finds a slightly smaller value

$$M_A = (1.001 \pm 0.020) \text{ GeV}. \quad (9)$$

The second method makes use of the so-called Adler-Gilman relation [9] which provides a chiral Ward identity establishing a connection between charged pion electroproduction at threshold and the isovector axial-vector current evaluated between single-nucleon states (see, e. g., [10, 11] for more details). At threshold (the spatial components of) the center-of-mass transition current for pion electroproduction can be written in terms of two s-wave amplitudes E_{0+} and L_{0+}

$$e\mathbf{M}|_{\text{thr}} = \frac{4\pi W}{m_N} [i\boldsymbol{\sigma}_\perp E_{0+}(k^2) + i\boldsymbol{\sigma}_\parallel L_{0+}(k^2)],$$

where W is the total center-of-mass energy, k^2 is the four momentum transfer squared of the virtual photon, and $\boldsymbol{\sigma}_\parallel = \boldsymbol{\sigma} \cdot \hat{k}$ and $\boldsymbol{\sigma}_\perp = \boldsymbol{\sigma} - \boldsymbol{\sigma}_\parallel$. The reaction $p(e, e'\pi^+)n$ has been measured at MAMI at an invariant mass of $W = 1125$ MeV (corresponding to a pion center-of-mass momentum of $|\mathbf{q}^*| = 112$ MeV) and photon four-momentum transfers of $-k^2 = 0.117, 0.195$ and 0.273 GeV^2 [12]. Using an effective-Lagrangian model an axial mass of

$$\bar{M}_A = (1.077 \pm 0.039) \text{ GeV}$$

was extracted, where the bar is used to distinguish the result from the neutrino scattering value. In the meantime, the experiment has been repeated including an additional value of $-k^2 = 0.058 \text{ GeV}^2$ [13] and is currently being analyzed. The global average from several pion electroproduction experiments is given by [5]

$$\bar{M}_A = (1.068 \pm 0.017) \text{ GeV}. \quad (10)$$




Fig. 1. Pion pole contribution to $\mathring{G}_P(q^2)$

It can be seen that the values of Eqs. (8) and (9) for the neutrino scattering experiments are smaller than Eq. (10) for the pion electroproduction experiments. The discrepancy was explained in heavy baryon chiral perturbation theory [14]. It was shown that at order $\mathcal{O}(q^3)$ pion loop contributions modify the k^2 dependence of the electric dipole amplitude E_{0+} from which M_A is extracted. These contributions result in a change of

$$\Delta M_A = 0.056 \text{ GeV}, \quad (11)$$

bringing the neutrino scattering and pion electroproduction results for the axial mass into agreement.

3 Induced pseudoscalar form factor $G_P(q^2)$

The induced pseudoscalar form factor $G_P(q^2)$ is less known than the axial form factor $G_A(q^2)$. A complete overview over the theoretical and experimental situation can be found in [6].

The axial-vector current is divergence-free in the chiral limit, $\partial_\mu A^{\mu,a} = 0$. In combination with the Dirac equation one obtains from Eq. (2) in the chiral limit that the equation

$$4\mathring{m}_N^2 \mathring{G}_A(q^2) + q^2 \mathring{G}_P(q^2) = 0 \quad (12)$$

must hold, where $\mathring{}$ denotes quantities in the chiral limit (see, e. g., [15] for a detailed discussion). Equation (12) has two possible solutions. The first solution is that $\mathring{m}_N = 0$ and $\mathring{G}_P(q^2) = 0$. However, the nucleon mass in the chiral limit does not vanish. The second solution is that $\mathring{G}_A(q^2)$ and $\mathring{G}_P(q^2)$ are related via

$$\mathring{G}_P(q^2) = -\frac{4\mathring{m}_N^2 \mathring{G}_A(q^2)}{q^2}. \quad (13)$$

Since $G_A(0) = g_A \approx 1.27$, it follows that $\mathring{G}_P(q^2)$ has a pole for $q^2 \rightarrow 0$. The behavior of $\mathring{G}_P(q^2)$ for $q^2 \rightarrow 0$ is interpreted as stemming from a pion pole contribution (see Fig. 1). The most general expression for the diagram of Fig. 1 in the chiral limit is given by

$$-\frac{4\mathring{m}_N F(q^2) \mathring{g}_{\pi N}(q^2)}{q^2 - \mathring{\Sigma}(q^2)} \bar{u}(p') \frac{q^\mu}{2\mathring{m}_N} \gamma_5 \frac{\tau^a}{2} u(p), \quad (14)$$

where $F(q^2)$ denotes the coupling of the axial source to the pion, $\overset{\circ}{g}_{\pi N}(q^2)$ is the pion nucleon coupling in the chiral limit and $\overset{\circ}{\Sigma}(q^2)$ stands for the pion self energy with $\overset{\circ}{\Sigma}(0) = \overset{\circ}{\Sigma}'(0) = 0$. Comparing with Eq. (2) one finds the pion pole diagram contribution

$$- \frac{4 \overset{\circ}{m}_N F(q^2) \overset{\circ}{g}_{\pi N}(q^2)}{q^2 - \overset{\circ}{\Sigma}'(q^2)} \quad (15)$$

to $\overset{\circ}{G}_P(q^2)$. The limit $q^2 \rightarrow 0$ is given by

$$\lim_{q^2 \rightarrow 0} q^2 \overset{\circ}{G}_{P,\pi\text{-pole}}(q^2) = -4 \overset{\circ}{m}_N F \overset{\circ}{g}_{\pi N} \quad (16)$$

which by comparison with Eq. (12) leads to the famous Goldberger-Treiman relation [16,17]

$$\frac{\overset{\circ}{g}_A}{F} = \frac{\overset{\circ}{g}_{\pi N}}{\overset{\circ}{m}_N}. \quad (17)$$

While the Goldberger-Treiman relation contains quantities in the chiral limit, it is interesting to note that it is satisfied to about 2% in the real world.

Information on $G_P(q^2)$ is mainly extracted from muon capture experiments. The induced pseudoscalar coupling g_P is defined as

$$g_P = \frac{m_\mu}{2m_N} G_P(q^2 = -0.88m_\mu^2). \quad (18)$$

It is mostly this quantity that has been determined by experiments, although one pion production experiment measured the q^2 dependence of $G_P(q^2)$ [18]. The weighted world average from ordinary muon capture (OMC),

$$\mu^- + p \rightarrow \nu_\mu + n, \quad (19)$$

is [5]

$$g_P = 8.79 \pm 1.92. \quad (20)$$

Most OMC experiments have used a liquid hydrogen target, which results in uncertainties due to the formation of $p\mu p$ molecules. For an exact determination of g_P the ortho-para transition rate has to be known. The current MuCap experiment at PSI [19] is using a hydrogen gas target and thereby avoiding these complications.

The pseudoscalar coupling g_P can also be determined from radiative muon capture (RMC),

$$\mu^- + p \rightarrow \nu_\mu + n + \gamma. \quad (21)$$

The value of g_P extracted from these experiments is [6]

$$g_P = 12.3 \pm 0.9 \quad (22)$$

and does not agree with the OMC result. However, a recent measurement at TRIUMF [20] found a new value for the ortho-para transition rate in the $p\mu p$ molecule of

$$A_{op}^{new} = (11.1 \pm 1.7 \pm_{0.6}^{0.9}) \times 10^4 s^{-1}, \quad (23)$$

which is significantly larger than the previous result [21] of

$$A_{op}^{old} = (4.1 \pm 1.4) \times 10^4 s^{-1}. \quad (24)$$

With the new value the RMC experiment gives [20]

$$g_P = 10.6 \pm 1.1, \quad (25)$$

while the average of the liquid hydrogen results is modified to be

$$g_P = 5.6 \pm 4.1. \quad (26)$$

Theoretically, $G_P(q^2)$ has been determined using heavy baryon chiral perturbation theory [22,23], which at order $\mathcal{O}(q^3)$ gives

$$g_P = 8.23. \quad (27)$$

4 $G_A(q^2)$ and $G_P(q^2)$ in Lorentz-invariant ChPT

With the introduction of renormalization schemes such as infrared regularization [24] or the extended-on-mass-shell scheme [25] the calculation of the isovector axial-vector current form factors in a manifestly Lorentz-invariant formulation of baryon chiral perturbation theory has been made possible [26].

The results of such a calculation up to and including order $\mathcal{O}(q^4)$ for $G_A(q^2)$ and $G_P(q^2)$ [27] are shown in Fig. 2 and Fig. 3, respectively. The analytic expression for $G_A(q^2)$ reads

$$G_A(q^2) = g_A + \frac{1}{6} g_A \langle r^2 \rangle_A q^2 + \frac{g_A^3}{4F^2} H(q^2), \quad (28)$$

where $\langle r^2 \rangle_A$ is the mean square axial radius and $H(q^2)$ contains contributions from loop diagrams with $H(0) = H'(0) = 0$. One can see that the contributions from $H(q^2)$ are small and the data is only described for very low values of momentum transfer. The calculation at order $\mathcal{O}(q^4)$ gives

$$g_P = 8.09 \quad (29)$$

and one can clearly see the pion pole contribution in Fig. 3.

The situation for the axial form factor $G_A(q^2)$ can be compared to the electromagnetic form factors, which in ChPT at order $\mathcal{O}(q^4)$ only describe the data for very low momentum transfers as well. In [28] a reformulation of infrared renormalization was used to include vector mesons as explicit degrees of freedom. This resulted in a better description of the data up to about $-q^2 \approx 0.4 \text{ GeV}^2$. Similar to the case of the electromagnetic form factors one can include axial-vector mesons in the formalism of baryon ChPT [27]. The Lagrangian for the coupling of the axial-vector meson to pions reads [29]

$$\mathcal{L}_{\pi A}^{(3)} = f_A \text{Tr}[A_{\mu\nu} F_-^{\mu\nu}], \quad (30)$$

while the coupling to the nucleon is given by

$$\mathcal{L}_{AN}^{(0)} = G_{AN} \bar{\Psi} A^\mu \gamma_\mu \gamma_5 \Psi. \quad (31)$$




Fig. 2. $G_A(q^2)$ in chiral perturbation theory at $\mathcal{O}(q^4)$ (solid line). The dashed line is a dipole fit. The data compilation is taken from [5].




Fig. 3. $G_P(q^2)$ in chiral perturbation theory at $\mathcal{O}(q^4)$.

The contributions of the diagrams containing the axial vector meson are

$$G_A^{AVM}(q^2) = -8f_A G_{AN} \frac{q^2}{q^2 - M_{a_1}^2} \quad (32)$$

to the axial form factor and

$$G_P^{AVM}(q^2) = 32m_N^2 f_A G_{AN} \frac{1}{q^2 - M_{a_1}^2} \quad (33)$$

to the induced pseudoscalar form factor, respectively. One sees that effectively only one new coupling constant, namely $f_A G_{AN}$, appears. This coupling constant can be fitted to the data [27].

5 Summary

A short overview over the experimental and theoretical situation of the nucleon form factors of the isovector axial-vector current was given. The axial form factor $G_A(q^2)$ has been determined by two types of experiments, neutrino scattering and pion electroproduction. The results from these two methods agree once pion loop corrections to the electroproduction amplitude have been taken into account. The situation for the induced pseudoscalar form factor $G_P(q^2)$ is less clear. The results obtained for the induced pseudoscalar coupling g_P from ordinary and radiative muon capture do not agree. However, a recent measurement of the ortho-para transition rate in $\mu\mu p$ molecules results in a significant change of the previous results for g_P . A reformulation of the infrared renormalization in baryon ChPT allows for the inclusion of axial-vector mesons in the calculation of the form factors, which could result in a better description of the experimental data. Only one new low-energy constant effectively appears, which can be fitted to the data.

This work was supported by the Deutsche Forschungsgemeinschaft (SFB 443).

References

1. H. y. Gao, Int. J. Mod. Phys. E **12** (2003) 1 [Erratum-ibid. E **12** (2003) 567].
2. J. Friedrich and Th. Walcher, Eur. Phys. J. A **17** (2003) 607.
3. C. E. Hyde-Wright and K. de Jager, Ann. Rev. Nucl. Part. Sci. **54** (2004) 217.
4. D. H. Wilkinson, Eur. Phys. J. A **7** (2000) 307.
5. V. Bernard, L. Elouadrhiri and U.-G. Meißner, J. Phys. G **28** (2002) R1.
6. T. Gorringer and H. W. Fearing, Rev. Mod. Phys. **76** (2004) 31.
7. W.-M. Yao et al., J. Phys. G **33**, (2006) 1.
8. H. Budd, A. Bodek and J. Arrington, arXiv:hep-ex/0308005.
9. S. L. Adler and F. J. Gilman, Phys. Rev. **152** (1966) 1460.
10. S. Scherer and J. H. Koch, Nucl. Phys. A **534** (1991) 461.
11. T. Fuchs and S. Scherer, Phys. Rev. C **68** (2003) 055501.
12. A. Liesenfeld *et al.* [A1 Collaboration], Phys. Lett. B **468** (1999) 20.
13. D. Baumann, PhD Thesis, Johannes Gutenberg-Universität, Mainz (2004).
14. V. Bernard, N. Kaiser and U.-G. Meißner, Phys. Rev. Lett. **69** (1992) 1877.
15. S. Scherer, in *Advances in Nuclear Physics, Vol. 27*, edited by J. W. Negele and E. W. Vogt (Kluwer Academic/Plenum, New York 2003) 277-538.
16. M. L. Goldberger and S. B. Treiman, Phys. Rev. **110** (1958) 1178.
17. M. L. Goldberger and S. B. Treiman, Phys. Rev. **111** (1958) 354.
18. S. Choi *et al.*, Phys. Rev. Lett. **71**, (1993) 3927.
19. MuCap Collaboration (Spokespersons P. Kammel and C. Petitjean), <http://www.npl.uiuc.edu/exp/mucapture/>.

- 20. J. H. D. Clark *et al.*, Phys. Rev. Lett. **96** (2006) 073401.
- 21. G. Bardin *et al.*, Phys. Lett. B **104** (1981) 320.
- 22. V. Bernard, N. Kaiser and U.-G. Meißner, Phys. Rev. D **50** (1994) 6899.
- 23. H. W. Fearing, R. Lewis, N. Mobed and S. Scherer, Phys. Rev. D **56** (1997) 1783.
- 24. T. Becher and H. Leutwyler, Eur. Phys. J. C **9** (1999) 643.
- 25. T. Fuchs, J. Gegelia, G. Japaridze and S. Scherer, Phys. Rev. D **68** (2003) 056005.
- 26. S. i. Ando and H. W. Fearing, arXiv:hep-ph/0608195.
- 27. M. R. Schindler, T. Fuchs and S. Scherer, in preparation.
- 28. M. R. Schindler, J. Gegelia and S. Scherer, Eur. Phys. J. A **26** (2005) 1.
- 29. G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, Phys. Lett. B **223** (1989) 425.